

# Please proofread this information

The following is a LaTeX to HTML translation of the abstract information you entered for the Solar Physics Division Meeting 2000. This partial translation is how your abstract will appear online. The LaTeX in your abstract will be fully translated in The Bulletin of the American Astronomical Society (BAAS).

Please proof it and press the COMMIT button on the bottom of this form.

---

## Apparent Associations Between the Solar Interior, Corona, and Solar Wind

*R Woo, J.W. Armstrong (JPL/Caltech), S.R. Habbal (CfA)*

Using polarized brightness (pB) measurements made by the High Altitude Observatory (HAO) Mauna Loa Mk III K-coronameter, we investigate the daily changes of path-integrated density at 1.15 Ro representing mainly longitudinal structure but also temporal variations near the Sun. We relate these coronal features to the interior of the Sun by comparing them with the subsurface large-scale velocity patterns associated with torsional oscillations. Specifically, we find that the latitudinal profile of the correlation of density variations between latitudes separated by 20 deg is similar to that of the subsurface zonal variations of the Sun's differential rotation from the same time period in 1996. From equator to pole, bands of high and low density correlation appear to be associated with those of slower (retrograde) and faster (prograde) rotation, respectively. We also show that polar coronal holes are distinguished by non-recurring longitudinal structure, as opposed to recurring structure in the neighboring quiet Sun, with the standard deviation of the longitudinal variations of pB being half of that of the quiet Sun. These distinct signatures of the longitudinal structure of coronal density including their dependence on latitude are also present in 1993-1994, and replicated in the Ulysses proton density measurements of the distant solar wind during the same time, showing that polar coronal holes extend radially into the solar wind. Since polar coronal holes and the quiet Sun are associated with retrograde rotation, and the polar coronal hole boundary with prograde rotation, the manifestation of these coronal features in the solar wind also reflect the association of the solar wind with the dynamics of the solar interior.

**Presentation Type:** oral

**Category:** 5. Outer atmosphere: Corona

**Submitter:** Richard Woo

**Member ID:** 15048

**Presenter email address:** richard.woo@jpl.nasa.gov

**Presenter phone:** 818 354-3945

**Correspondent address:** MS 238-725 Jet Propulsion Laboratory 4800 Oak Grove Drive  
Pasadena, CA 91109

**New address flag:** yes

---

If the information is correct, press the "Commit" button below. If the information is *not* correct, please use the **Back** button on your browser to return to the input form and correct the problem.

When you are satisfied with your submission, please print this page for future reference.

**Commit**

Thank you.